

# Aspects of the Davenport Constant for Finite Abelian Groups

Anamitro Biswas  
anamitroappu@gmail.com

[joint work with Dr. Eshita Mazumdar]

18th International Conference of IMBIC  
Mathematical Sciences for Advancement of Science and Technology  
(MSAST) 2024  
Dec 21st-23rd

A sequence, also called a *multiset*, is a member of the free abelian group  $\mathcal{F}(G)$  generated by  $G$ . We shall denote by

- *juxtaposition* when elements form a **sequence**, and by
- *addition* the **group operation**.

Zero-sum problems in additive number theory:

- 1 Conditions which ensure that given sequences have non-empty zero-sum subsequences with prescribed properties.
- 2 Structure of extremal sequences which have no zero-sum subsequences.

Baayen, Erdős and Davenport posed the problem to determine

$$D(G) = \min \{|S| : S \in \mathcal{F}(G) \text{ has a non-trivial zero subsum}\}$$

called the *Davenport constant* for group  $G$ .



$D(G) \leq |G|$  for any group  $G$ .


*Proof.* Let  $S = x_1 x_2 \dots x_n \in \mathcal{F}(G)$  where  $|G| = n$ . Consider

$$s_1 = x_1$$

$$s_2 = x_1 + x_2$$

$$\vdots$$

$$s_n = x_1 + x_2 + \dots + x_n$$



If all the  $s_i$ 's are distinct, we must have  $0 \in \{s_1, \dots, s_n\}$  since  $G$  contains only  $n$  elements. Else, we have by the Pigeon-Hole Principle,

$\exists i \neq j \ni s_i = s_j$ . Then,  $x_{i+1} \dots x_j$  is a zero-sum subsequence. So,  $D(G) \leq |G|$ . □

Particularly for  $G = C_n = \langle 1 \rangle$  (the cyclic group of order  $n$ ) we can construct the sequence  $S = \underbrace{11 \dots 1}_{n-1}$  such that  $0 \notin [S]$  (the set of subsums of sequence  $S$  including  $\sigma(S)$ ). Thus  $D(C_n) \geq n \Rightarrow D(C_n) = n$ .

## Theorem (Olson, 1961)

$$D\left(\bigoplus_{i=1}^d C_{p^{e_i}}\right) = 1 + \sum_{i=1}^d (p^{e_i} - 1)$$

Olson further conjectured that for any finite abelian group

## Conjecture (Olson, 1961)

$$D\left(\bigoplus_{i=1}^d C_{n_i}\right) = 1 + \sum_{i=1}^d (n_i - 1) = D^*(G) \text{ where } n_i \mid n_{i+1} \text{ for } i \in \{1, \dots, d-1\}.$$

Too good to be true, this conjecture isn't in fact, since, for example, we have

## Geroldinger and Schneider, 1992

For odd  $m, n \ni 3 \leq m|n$ ,

$$D\left(C_m \oplus C_n^2 \oplus C_{2n}\right) > D^*\left(C_m \oplus C_n^2 \oplus C_{2n}\right).$$

However, we have  $D(C_m \oplus C_n) = D^*(C_m \oplus C_n)$  where  $m \mid n$  and all known counterexamples are of rank  $\geq 4$ . So, apart from characterization of the groups for which this conjecture holds, it is also an open problem whether the conjecture holds for groups of rank 3.

Theorem (Bhowmik and Schlage-Puchta)

$$D(C_3 \oplus C_3 \oplus C_{3d}) = D^*(C_3 \oplus C_3 \oplus C_{3d}) \quad \forall d \in \mathbb{N}.$$

Conjecture

Fixed  $5 \leq p \in \mathbb{P}$ ,  $G \cong C_p^3 \oplus C_2$ .  
 $S = (x_1, y_1) \dots (x_{4p-2}, y_{4p-2}) \in \mathcal{F}(G) \ni$

$$y_1 = \dots = y_r = 1$$

$$y_{r+1} = \dots = y_{4p-2} = 0$$

for even  $r \in [2p+2, 4p-6]$ . Then  $0 \in [S]$ .

Theorem (Sheikh, 2017)

$$D(C_p \oplus C_p \oplus C_{2p}) = D^*(C_p \oplus C_p \oplus C_{2p})$$

for all those primes  $p$  for which the previous Conjecture holds.

Theorem (Sheikh, 2017)

$$D(C_5 \oplus C_5 \oplus C_{5d}) \leq D^*(C_5 \oplus C_5 \oplus C_{5d}) + 4 \quad \forall d \in \mathbb{N}.$$

With the help of computer, Sheikh further confirmed,  $D(C_5 \oplus C_5 \oplus C_{10}) = D^*(C_5 \oplus C_5 \oplus C_{10}) = 18$ . Generalizing Sheikh's approach,

## Conjecture

Fix  $p, q$  ( $p \neq q$ ),  $p \in \mathbb{P}$ ; define  $G := C_p^d \oplus C_q$ .

Let  $m = p(q+2) - 2$ .

Let  $S = (x_1, y_1) \dots (x_m, y_m) \in \mathcal{F}(C_p^d \oplus C_q)$ . Suppose

$$y_{\sum_{i=1}^t r_i+1} = \dots = y_{\sum_{i=1}^{t+1} r_i} = t+1 \quad (t \in [0, q-1])$$

where  $r = \sum_{i=1}^{q-1} r_i$ . If

①  $r \in [pq+1, p(q+2)-2]$  and

②  $\sum_{i=1}^{q-1} ir_i \equiv 0 \pmod{q}$ ,

then  $0 \in [S]$ .

## Theorem (me and Eshita Ma'am)

Let  $p$  be a prime such that the property stated in the Conjecture holds. Then,  $D(C_p^d \oplus C_q) = D^*(C_p^d \oplus C_q)$ .

(Bhowmik and Schlage-Puchta, 2007) For  $G \cong C_3 \oplus C_3 \oplus C_{3q}$ ,  $q \in \mathbb{N}$ ,  $D(G) = D^*(G)$ . So the Conjecture is true for  $p = 3, d = 3$  at least.

For the group  $C_5 \oplus C_5 \oplus C_{15}$ , the conjecture can be further reduced to the assumption for only the cases where  $10 \leq r_2 - \lfloor \frac{r_2 - r_1}{3} \rfloor$ , by constructing subsequences of this form:

$$S' = \left( \prod_{i=1}^{\alpha} (x_{r+i}, y_{r+i}) \right) \left( \prod_{i=1}^{r_1} \sigma((x_i, y_i) (x_{r_1+i}, y_{r_1+i})) \right) \left( \prod_{i=\frac{r_1+\ell}{3}}^{\lfloor \frac{r_2}{3} \rfloor} \sigma((x_{r_1+i}, y_{r_1+i}) (x_{r_1+i+1}, y_{r_1+i+1}) (x_{r_1+i+2}, y_{r_1+i+2})) \right)$$

where  $\ell = 3 \lfloor \frac{r_1}{3} \rfloor$ ,  $\alpha = s - r = 23 - r$  and  $\beta = \min(r_1, 23 - \alpha - r_1) = r_1$ .

## the $r$ -wise Davenport constant

There can be a number of possible ways to generalize the idea of Davenport constant which just requires *one* sequence that adds to zero.

### Definition (Girard and Schmid)

For  $r \in \mathbb{N}$ , the  $r$ -wise Davenport constant of group  $G$ ,

$$D_r(G) = \min \{k \in \mathbb{N} \mid S \in \mathcal{F}(G) \text{ \& } |S| \geq k \\ \Rightarrow S \text{ has } r \text{ disjoint zero-sum subsequences}\}.$$

It is clear that  $D_r(G) \leq D_{r+1}(G)$ .

### Theorem (Girard and Schmid, 2019)

- If  $n, r \in \mathbb{N}$ ,  $D_r(C_n) = rn$ .
- Let  $G \cong C_m \oplus C_n$  where  $m|n$ . Then,  $D_r(G) = rn + m - 1$ .



Let  $p$  be an odd prime and  $G \cong C_{p^{e_1}} \oplus C_{p^{e_2}} \oplus \cdots \oplus C_{p^{e_d}} \ni e_i \leq e_{i+1}$  for  $i \in [1, d-1]$ .  $S = \underbrace{(1, 0, \dots, 0)}_d {}^{rp^{e_d}-1} (0, 1, \dots, 0) {}^{p^{e_d-1}-1} \dots$

$(0, 0, \dots, 0, 1) {}^{p^{e_1}-1}$  does not have  $r$  disjoint zero-sum subsequences.

Therefore,  $D_r(G) \geq rp^{e_d} + \sum_{i=1}^{d-1} p^{e_i} - d + 1$ .

$\eta_r(G) = \min \{k \in \mathbb{N} \mid \forall S \in \mathcal{F}(G) \ni |S| \geq k, (0_{\text{small}})^r \in [S]\}$ . By  $0_{\text{small}}$  we mean a small zero subsum, i.e., a subsum of length  $\leq \exp(G)$ .

Theorem (Fan, Gao, Wang, Zhong (2013))

$m \in \mathbb{N}$ ; let  $H$  be a finite abelian group  $\ni$

- $\exp(H) \mid m$
- $m \geq D(H)$
- $D(C_m \oplus C_m \oplus H) = 2m + D(H) - 2$

Then  $\eta_1(C_m \oplus H) \leq 2m + D(H) - 2$ .

If  $p^{e_d} \geq 1 + \sum_{i=1}^{d-1} (p^{e_i} - 1)$ , then  $\eta_1(G) \leq D(G) + \exp(G)$ . Let  $1 \leq r \in \mathbb{Z}$ .

If  $\eta_1(G) \leq D(G) + \exp(G)$ , then  $\eta_r(G) \leq D(G) + r \exp(G)$ . For such a group,  $D_r(G) \leq \eta_r(G) = D(G) + (r - 1) \exp(G)$ .

Theorem (Geroldinger and Halter-Koch, 2006)

Let  $p$  be an odd prime and  $G \cong C_{p^{e_1}} \oplus C_{p^{e_2}} \oplus \cdots \oplus C_{p^{e_d}} \ni e_i \leq e_{i+1}$  for  $i \in [1, d - 1]$ . Then,  $D_r(G) = rp^{e_d} + \sum_{i=1}^{d-1} p^{e_i} - d + 1$ .

Theorem (Delorme, Ordaz and Quiroz (2001))

$p \in \mathbb{P}, n \geq 2$  &  $(m, p^n) = 1$ . Then, for  $G \cong C_p \oplus C_p \oplus C_{p^n m}$ ,  $D(G) = D^*(G)$ .

Delorme, Ordaz and Quiroz (2001)

Let  $H \triangleleft G$  and  $r \in \mathbb{N}$ , then  $D_r(G) \leq D_{D_r(H)}(G/H)$  and  $D(G) \geq D(H) + D(G/H) - 1$ .

## Theorem (Me and Eshita Ma'am)

$$G \cong C_{n_1} \oplus C_{n_2} \oplus \cdots \oplus C_{n_d} \quad (n_1, n_2, \dots, n_d \in \mathbb{N})$$

$$\text{i.e., } G = C_{\prod_{j=1}^{\ell} p_j^{e_1^{(j)}}} \oplus C_{\prod_{j=1}^{\ell} p_j^{e_2^{(j)}}} \oplus \cdots \oplus C_{\prod_{j=1}^{\ell} p_j^{e_d^{(j)}}} \quad \text{where}$$

- w.l.o.g.  $e_i^{(j)} \in \mathbb{Z} \ni 0 \leq e_i^{(j)} \leq e_{i+1}^{(j)} \quad \forall 1 \leq j \leq \ell$
- but all  $e_i^{(j)}$ 's are not zero for each  $j \in \{1, \dots, \ell\}$  ( $1 \leq i \leq d$ )
- $p_1, p_2, \dots, p_{\ell}$  primes  $\ni p_j^{e_d^{(j)}} \geq 1 + \sum_{i=1}^{d-1} (p_j^{e_i^{(j)}} - 1) \quad \forall j = 1, 2, \dots, \ell.$

$$\text{Let } \varphi(p_j) = \sum_{i=1}^{d-1} p_j^{e_i^{(j)}} - d + 1 \text{ for } j = 1, \dots, \ell.$$

$$\begin{aligned} \text{Then, } r \prod_{j=1}^{\ell} p_j^{e_d^{(j)}} + \sum_{i=1}^{d-1} \left( \prod_{j=1}^{\ell} p_j^{e_i^{(j)}} - 1 \right) &\leq D_r(G) \\ &\leq r \prod_{j=1}^{\ell} p_j^{e_d^{(j)}} + \sum_{m=1}^{\ell-1} \left( \left( \prod_{j=m+1}^{\ell} p_j^{e_d^{(j)}} \right) \varphi(p_m) \right) + \varphi(p_{\ell}). \end{aligned}$$

## Corollary (a generalization of Delorme et al)

For  $G = C_{p^{e_1}} \oplus C_{p^{e_2}} \oplus \cdots \oplus C_{p^{e_{d-1}}} \oplus C_{rmp^{e_d}}$  with  $e_j \leq e_{j+1}$  such that

$$p^{e_d} \geq 1 + \sum_{i=1}^{d-1} (p^{e_i} - 1),$$

$$D_r(G) = rmp^{e_d} + \sum_{i=1}^{d-1} p^{e_i} - d + 1.$$

## Result

The error becomes negligible, i.e.,  $\frac{\text{upper bound}}{\text{lower bound}} \rightarrow 1$  if

- either  $p_j$ 's [ $j \in \{1, \dots, \ell\}$ ] are large;
- $e_i^{(j)}$  ( $j = 1, \dots, \ell; i = 1, \dots, d$ )'s are higher natural numbers;
- $r$  increases.

$$\text{error} := \frac{\text{upper bound} - \text{lower bound}}{\text{lower bound}}$$

[error has been throughout multiplied by 100 for easy visualization]

Group		$r = 1$	$r = 2$	$r = 3$	$r = 4$	$r = 5$
$C_{2.3.5} \oplus C_{2^2.3^2.5^2} \oplus C_{2^3.3^3.5^3}$	UB	41778	68778	95778	122778	149778
	LB	27928	54928	81928	108928	135928
	diff	13850				
	err	49.59181	25.21483	16.90509	12.71482	10.18922
$C_{3.5.7} \oplus C_{3^2.5^2.7^2} \oplus C_{3^3.5^3.7^3}$	UB	1596033	2753658	3911283	5068908	6226533
	LB	1168753	2326378	3484003	4.641628	5799253
	diff	427280				
	err	36.55862	18.36675	12.26405	9.205391	7.367845
$C_{5.7.11} \oplus C_{5^2.7^2.11^2} \oplus C_{5^3.7^3.11^3}$	UB	69921553	126988178	184054803	241121428	298188053
	LB	57215233	114281858	171348483	228415108	285481733
	diff	12706320				
	err	22.207932	11.118405	7.415484	5.562819	4.450835

Group

$$C_{31^{1.47}.101} \oplus C_{31^{2.47^2}.101^2} \oplus C_{31^{3.47^3}.101^3}$$

	$r = 1$	$r = 5$
UB	3292613286703417	16039460139018988
LB	3186733368408697	15933580220724268
diff	105879918294720	
err	3.3225220	0.6645080

$$C_{31^{12.47^3}.101^3} \oplus C_{31^{18.47^9}.101^5} \oplus C_{31^{17.47^{21}}.101^7}$$

$r = 1$

Upper bound=314378927707039117076594641960472205699918246393796134855347548904541388800

Lower bound= 314378927707027215691704348813743973043171844658271204789424845095104937984

Difference= 11890447077876842522672189603253739181827333079849168416014336

Error= 0.0000000000037822023

$r = 5$

Upper bound= 1571894638535147728759342621748217544441028696720224309918829150723580624896

Lower bound=1571894638535135877591266211694935422470597680646030708671750002626420277248

Error= 0.000000000000756440

# References



G. Bhowmik and J. Schlage-Puchta, Davenport's constant for groups of the form  $\mathbb{Z}_3 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_{3d}$ . In Additive combinatorics, volume 43 of CRM Proc. Lecture Notes, 307-326. *Amer. Math. Soc.*, Providence, RI, (2007).



A. Biswas and E. Mazumdar, *Davenport constant for finite abelian groups with higher rank*, arXiv:2402.09999 (2023).



A. Geroldinger and F. Halter-Koch, Non-Unique Factorizations: Algebraic, Combinatorial and Analytic Theory, *Chapman & Hall/CRC* (2006).



A. Geroldinger, R. Schneider, On Davenport's constant, *J. Combin. Theory Ser. A* 61, 147—152, (1992).



B. Girard and W. A. Schmid, Direct zero-sum problems for certain groups of rank three, *J. Number Theory*, 197, 297—316, (2019).



J. E. Olson, A Combinatorial Problem on Finite Abelian Groups, I, *J. Number Theory*, 1, 8–10, (1969).



K. Rogers, A Combinatorial problem in Abelian groups, *Proc. Cambridge Phil. Soc.* 59, 559-562, (1963).



A. Sheikh, The Davenport constant of finite abelian groups, *Thesis, University of London*, (2017).



Thank you :)  
A handout for this talk can  
be found at:

[https://anamitro.  
github.io/files/  
anamitro\\_msast24.pdf](https://anamitro.github.io/files/anamitro_msast24.pdf)